

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
 Organization
 International Bureau



(43) International Publication Date
 4 March 2004 (04.03.2004)

PCT

(10) International Publication Number
WO 2004/017739 A1

- (51) International Patent Classification⁷: **A01N 65/00**
- (21) International Application Number:
 PCT/IT2003/000514
- (22) International Filing Date: 22 August 2003 (22.08.2003)
- (25) Filing Language: Italian
- (26) Publication Language: English
- (30) Priority Data:
 B02002A 000544 23 August 2002 (23.08.2002) IT
- (71) Applicant (for all designated States except US): **CERE-ALTOSCANA S.p.A.** [IT/IT]; Via Pistoiese, 43, I-50054 Fucecchio (IT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LAZZERI, Luca** [IT/IT]; Via Richa Giuseppe, 80, I-50100 Firenze (IT). **LEONI, Onofrio** [IT/IT]; Via Enrico Berlinguer, 125, I-40024 Castel San Pietro Terme (IT). **MANICI, Luisa, M.** [IT/IT]; Via Bottego, 28, I-40131 Bologna (IT). **PALMIERI, Sandro** [IT/IT]; Via Molino di Pescarola, 72, I-40131 Bologna (IT). **PATALANO, Giampiero** [IT/IT]; Via A. Batteli, 49, I-56100 Pisa (IT).
- (74) Agents: **JORIO, Paolo et al.**; C/O Studio Torta S.R.L., Via Viotti, 9, 10121 Torino (IT).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
 — with international search report
 — before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: USE OF SEED FLOUR AS SOIL PESTICIDE

(57) Abstract: Use of vegetable seed flour, in particular Brassicaceae flour, as a cytotoxic agent with improving action.

WO 2004/017739 A1

5

USE OF SEED FLOUR AS SOIL PESTICIDE

10

TECHNICAL FIELD

The present invention concerns a use of seed flour as a soil improving agent, as a cytotoxic agent for
15 soil pathogens and parasites, and as a fungitoxic agent for soil fungal pathogens.

BACKGROUND ART

In the agricultural sector methyl bromide is commonly used to limit the growth of fungal pathogens,
20 nematodes, wireworms, insects, bacteria and weeds. However, methyl bromide has a relatively high environmental impact: reacting with the ozone present in the upper layer of the atmosphere, it transforms it into Br_2O , contributing to destruction of the ozone.

25 Recently, to replace methyl bromide, molecules of vegetable origin with cytotoxic activity for soil pathogens and parasites have been proposed. Among these molecules, derivatives of glucosinolates (GLs)

- 2 -

generated by hydrolysis catalysed by the enzyme myrosinase (MYR) or by thermolysis have been proposed. The GLs are anionic thioglucosidic compounds present in variable quantities and ratios in a number of organisms
5 and vegetable tissues. The molecular structure of the approximately 120 GLs currently isolated and characterised consists of a common functional group and a lateral chain which can be of aliphatic, aromatic or heteroaromatic type.

10 The GLs, in the presence of the enzyme MYR (thioglucoside glucohydrolase, EC 3.2.3.1), are hydrolysed with formation of β -D-glucose, sulphate ion and a number of derivatives such as isothiocyanates, nitriles or thiocyanates (see, for example, Tookey H.
15 L., Van Etten C. H., Daxembichler M. E., *Toxic Constituent of Plant Foodstuffs*, edited by I. E. Liner, 1980, II edition, 4, 103-142).

Some of the above-mentioned derivatives of the GLs, in particular the isothiocyanates, have shown in vitro
20 cytotoxic properties towards nematodes (see Lazzeri L., Tacconi R., Palmieri S., *In Vitro Activity of Some Glucosinolates and Their Reaction Products toward a Population of the Nematode *Heterodera schachtii**, *J. Agric. and Food Chem.* 1993, 41, 825-829) and plant
25 fungal pathogens present in the soil (see, for example, Manici L.M., Leoni O., Lazzeri L., Galletti S.,

- 3 -

Palmieri S., Fungitoxic Activity of Some Glucosinolate Enzyme Derived Products Against the Main Soil-Borne Pathogens, *Pesticide Science*, 1999, 55, 486-488; and Manici L.M., Lazzeri L., Palmieri S., In Vitro
5 Fungitoxic Activity of Some Glucosinolates and their Enzyme-derived Products toward Plant Pathogenic Fungi, *J. Agric. and Food Chem.* 1997, 45, 2768-2773).

To obtain and use the above-mentioned derivatives of the GLs, extraction and/or production procedures are
10 necessary, which are relatively complex and costly and can also cause environmental problems (for example disposal of the by-products and/or the solvents used). Furthermore, generally some important derivatives of the GLs (for example allylisothiocyanate) are
15 relatively difficult to handle as they have a relatively high volatility. To overcome these negative aspects the technique of green manure of plants containing both GLs and the enzyme MYR (i.e. containing the GLs-MYR system) has been proposed.

20 The green manure technique of plants containing the GLs-MYR system has some negative aspects, however, typical of the green manure technique in general. In particular, the green manure technique requires cultivation operations that determine relatively high
25 labour costs and causes the loss of at least one growing cycle, with consequent damage for the farmer

- 4 -

due to loss of income.

DISCLOSURE OF INVENTION

The aim of the present invention is to provide a use of a seed flour as a soil improver in order to
5 reduce the above-mentioned disadvantages in a relatively simple and inexpensive way.

The present invention provides a use of a seed flour containing at least one glucosinolate and at least one enzyme chosen from the group consisting of:

- 10 - glucosidasic enzymes, and
 - thioglucosidasic enzymes;
 as a soil improver.

A further aim of the present invention is to provide a use of a seed flour as a cytotoxic agent for
15 soil pathogens and parasites.

The present invention provides a use of a seed flour containing at least one glucosinolate and at least one enzyme chosen from the group consisting of:

- 20 - glucosidasic enzymes, and
 - thioglucosidasic enzymes;
 as a cytotoxic agent for soil pathogens and soil parasites.

A further aim of the present invention is to provide a use of a seed flour as a fungitoxic agent.

25 The present invention provides a use of a seed flour containing at least one GL and at least one

- 5 -

enzyme chosen from the group consisting of:

- glucosidasic enzymes, and
- thioglucosidasic enzymes;

as a fungitoxic agent for fungal pathogens of the
5 soil.

Further characteristics of the present invention will be better clarified by the following non-restrictive description.

BEST MODE FOR CARRYING OUT THE INVENTION

10 Seeds containing GLs and at least one glucosidasic or thioglucosidasic enzyme are ground in order to obtain a fine uniform flour. Preferably the seeds contain the enzyme MYR, more preferably they belong to plants in the order of the Capparales and to the
15 families of the Akaniaceae, Bataceae, and even more preferably they are seeds of *Brassicaceae*, *Capparaceae*, *Caricaceae*, *Gyrostemonaceae*, *Limnanthaceae*, *Moringaceae*, *Pentadiplandraceae*, *Koeberliniaceae*, *Resedaceae*, *Salvodoraceae*, *Tropeolaceae* and/or
20 *Toviaraceae*.

Preferably the seeds used are seeds of *Brassicaceae* and/or *Capparaceae*, in particular they are seeds of *Brassica carinata* A. B., seeds of *Eruca Sativa* M., and/or seeds of *Barbarea verna*. Furthermore, it is
25 important to point out that it is possible to identify other vegetable varieties, the seeds of which contain

- 6 -

GLs and a glucosidasic or thioglucosidasic enzyme, analysing the seeds by means of known methods.

The seed flour thus obtained has a variable content of GLs, glucosidasic or thioglucosidasic enzyme, 5 proteinic material and oil according to the vegetable varieties and/or the ambient growing conditions. Furthermore, it is important to underline that when the vegetable varieties used vary, the related quantities of the different types of GLs contained in the seeds 10 can also vary.

For each vegetable variety the best growing techniques can be identified (i.e. by optimising, for example, the sowing period, density, type of soil and/or fertilising agents) in order to obtain a higher 15 content of GLs and glucosidasic or thioglucosidasic enzyme in the seeds.

It should also be noted that, preferably, to further increase the content of GLs and glucosidasic or thioglucosidasic enzyme, the above-mentioned seed flour 20 is de-oiled, in other words it is treated in order to extract at least part of the oil contained in it. The extraction is performed preferably at a relatively low temperature, preferably below 75°C, in particular at ambient temperature, so that the glucosidasic or 25 thioglucosidasic enzyme is not thermally deteriorated and, also after the oil extraction phase, it is still

- 7 -

able to perform its enzymatic activity. In this regard it is important to underline that the oil extracted can be used, for example, as a renewable and ecologically sustainable lubricant or hydraulic fluid.

5 The flour, prepared in the form of pellets, can be distributed over the soil relatively easily and at a relatively low cost. In the presence of water, derivatives of the GLs generated by hydrolysis catalysed by the glucosidasic or thioglucosidasic
10 enzyme develop in the soil. In this regard, it is important to underline that in order to hydrolyse the GLs, water is required, which is a reagent essential for triggering and developing the biochemical reaction.

For this reason, the non-hydrated flour has a
15 relatively high stability, can be fairly easily transported, stored for relatively long periods and used at any time, regardless of the ambient conditions.

Once the flour has been distributed on the soil and hydrated, the derivatives of the GLs generated by
20 enzymatic hydrolysis are able to perform their cytotoxic action even though the flour contains approximately 50% protein. Therefore, although it may be feared that the isocyanates generated by hydrolysis of the GLs, catalysed by the glucosidasic enzymes, or
25 more specifically by the thioglucosidasic enzyme MYR, may spontaneously interact with the free NH₂ groups of

the lysines and arginines contained in the proteins, to produce adducts of a different type that can be referred to derivatives of thiourea, the hydrolysis products of the free GLs are surprisingly still able to perform a relatively high improving action in the soil. The hydrolysis products of the GLs perform a fungitoxic action, in particular on fungal pathogens of the soil belonging to different taxonomic classes (for example Oomycetes: *Pythium* and *Phytophthora*; Sterile fungi: *Sclerotium rolfsii* and *Rhizoctonia solani*; Hyphomycetes: *Fusarium* spp and *Alternaria* spp.) and a cytotoxic action on soil pathogens and parasites, in particular nematodes (in particular *Meloidogyne incognita*), insects and weeds. Among the insects on which cytotoxic action is performed, the cytotoxic activity on wireworms is particularly interesting (for example *Agriotes sordidus*, *Agriotes utulatus* and *Agriotes brevis*). The fungitoxic activity on fungi in the genus *Fusarium* is also particularly interesting.

It is important to point out that use of the flour permits exploitation of concentrations of GLs substantially impossible to achieve with the green manure technique.

Note that it is possible to use either flour of seeds of one single vegetable variety, or a mixture of flour of seeds of a number of vegetable varieties. In

- 9 -

particular, the varieties used for their seeds can be chosen so as to create flour formulates with different modes of action, according to the need for persistence or volatility of the derivatives of the GLs and the required cytotoxic action. In fact, both characteristics (volatility and cytotoxic action) can vary according to the type of pathogen or parasite and the phase it is in at the time: vegetation (like mycelium) or conservation (sclerotium, oospore, chlamydospore or hibernating mycelium). Therefore, according to the modes of application (times and types of culture) of the target pathogen(s) and/or parasite(s) and the need for improvement, the compositions of the formulates can be varied. In this way it is also possible to obtain formulates for the control of telluric pathogens and parasites. Furthermore, it is possible to use the flour described above also in combination with other compounds, which in their turn can be of either synthetic or natural origin.

Further characteristics of the present invention will become clear from the description of some merely illustrative and non-restrictive examples. In particular, although the following examples refer to the use of seed flour of *Brassica carinata* A.B, *Eruca sativa* M. and *Barbarea verna* as fungitoxic agents, the

- 10 -

types of flour that can be used for said purpose are all those containing GLs and a glucosidasic or thioglucosidasic enzyme (in particular MYR). Furthermore, the flour containing GLs and a
5 glucosidasic or thioglucosidasic enzyme (in particular MYR) can be used also as a cytotoxic agent for other soil pathogens and parasites.

It is important to point out that the results illustrated in the following examples can vary in
10 quantitative but not qualitative terms, according to the vegetable varieties and ambient conditions in which the seeds are produced.

Example 1

This example describes a method for obtaining seed
15 flour of *Brassica carinata* A.B, *Eruca sativa* M. and *Barbarea verna*.

Seeds of *Brassica carinata* A.B, *Eruca sativa* M. and *Barbarea verna* are cleaned and then ground separately with an ordinary rolling machine until a fine uniform
20 flour is obtained. Each type of flour is separately de-oiled with *n*-hexane (Carlo Erba®) in a concentration of 1:10 (weight/volume), leaving the suspension to be stirred in an agitator for 12 hours at ambient temperature. Each type of flour is then separately
25 filtered on paper (Omniafiltra Superverloxx filter, Cartiera del Torano SpA, Naples), washed repeatedly

- 11 -

again with n-hexane and placed in a ventilated environment at ambient temperature for 24 hours in order to eliminate all remains of solvent.

Example 2

5 This example describes an analysis method to determine the content of GLs and the mirosinasic activity in seed flour.

Each type of flour, obtained as described in example 1, was analysed to determine the content in
10 GLs, using the procedure established by the regulation of the European Union 1864/90 for the analysis of rape seed (Official Journal of the European Communities, L 170, 03/07/1990, p. 0027-0034). The analysis was performed with HPLC technique (High Performance Liquid
15 Chromatography), using a Hewlett Packard® Chromatograph Mod. 1090L with diode detector ($\lambda = 229$ nm) and a column HP ODS Hypersil C18, 5 μ m 200 x 4.6 mm. The mobile phase consisted of a mixture of water and acetonitrile with a gradient from 1% to 22% of
20 acetonitrile in 22 minutes and with a flow of 2 lm/min. The temperature of the column was 35°C.

Furthermore the total myrosinase activity in the flour was also quantified via the pH-stat assay (Palmieri S., Iori R., Leoni, O. Comparison of Methods
25 for Determining Myrosinase Activity, J. Agric. Food Chem. 1987, 35, 617-621). The analyses were repeated

- 12 -

three times, giving the results in Table I, after calculation of the means and standard deviations.

Table I

Flour	GLs content ($\mu\text{moli g}^{-1}$ TQ)	Prevalent GL	Myrosinase activity (U g^{-1})
<i>B. carinata</i>	141.6 ± 4.8	Allyl GL (Sinigrine)	34.7 ± 0.4
<i>E. sativa</i>	161.1 ± 2.1	4- Methylthiobutyl GL (Glucorucin)	53.3 ± 4.2
<i>B. verna</i>	$143,7 \pm 0.6$	Fenyl-ethyl GL (Nasturtin)	2.4 ± 0.5

5 The data given in Table I indicate that after cold grinding and extraction with *n*-hexane, the GLs and the enzyme MYR in active form are present in the flour. Therefore the de-oiled seed flour of *P Brassica carinata* A.B, *Eruca sativa* M. and *Barbarea verna*
 10 contain the GLs-MYR system, and hence, in the presence of water, are able to produce derivatives of GLs generated by enzymatic hydrolysis via MIR.

The analysis method described above can be used to identify the vegetable varieties in the seeds of which
 15 the GLs-MYR system is present.

Example 3

This example describes a method for assessing the fungitoxic activity.

The fungitoxic activity test was performed
5 starting from an isolate of *Fusarium culmorum*, isolated
and identified as in Manici M.L. and Cerato C. (1992) -
Studio su alcuni funghi agenti di marciume dei tuberi
di patata; Informatore Fitopatologico 9: 41-46, and
references quoted in it. This isolate is catalogued as
10 fungo test at the micotech of the Experimental
Institute for Industrial Cultivation (Bologna) and was
chosen for the stability and regularity of growth that
characterise it.

The test provided for assessment of the toxicity
15 of the volatile compounds released by the flour
following hydration.

Doses of 0.01 g, 0.025 g, 0.05 g, 0.25 g and 1 g
of the seed flour of *Brassica carinata*, *Eruca sativa*
and *Barbarea verna* analysed in example 2 were tested
20 separately. The control, included for assessment of the
fungitoxic activity, consisted of pellets of commercial
organic soil improver based on fowl dung and torrefied
leather.

The assay was based on assessment of the reduction
25 of growth in colonies of *Fusarium culmorum* in the
presence of the products released by the flour

- 14 -

following hydration, with respect to growth of the same fungus in the presence of the control.

Colony discs of 4 mm taken from a one-week old colony of *Fusarium culmorum* were inoculated on a bottom of a Petri dish on an agar nutritive substrate (Potato Dextrose Agar) and immediately after overturned onto a bottom of a Petri dish containing the control and, separately, the flour in the different amounts (in a total volume of approximately 174 cm³) hydrated immediately beforehand. The two Petri dish bottoms joined as above were sealed with parafilm, and the resulting dishes incubated for the time necessary for the colony on the control to develop and invade at least $\frac{3}{4}$ of the dish. At this point the diameters of the colonies grown in the presence of the vapours released at the different doses of flour and control were measured.

The fungitoxic activity of the flour was expressed as inhibition effectiveness (EI) as a percentage with respect to the control, using the formula: $EI (\%) = \frac{[(\text{Diameter of Control} - \text{Diameter of Treated Sample}) / \text{Diameter of Control}] \times 100}{}$

Table II shows the results of the test.

The test was performed with three repetitions per treatment, therefore the data shown in table II are the means of three values.

- 15 -

Table II

Treatment	Dose	EI %
De-oiled flour	(g/dish)	
<i>Eruca sativa</i>	0.01	81
	0.025	100
	0.05	100
	0.25	100
	1	100
<i>Brassica carinata</i>	0.01	72
	0.025	100
	0.05	100
	0.25	100
	1	100
<i>Barbarea verna</i>	0.01	40
	0.05	100
	0.25	100
	1	100

The above data indicate that the products released by the seed flour of *Brassica carinata* A. B., *Eruca sativa*
 5 *M.* and *Barbarea verna*, following hydration, which activates the GLs-MYR system present in it, inhibit to a large extent growth of the fungus *Fusarium culmorum*.

Example 4

This example describes a method for assessing the
 10 nematocide activity on nematodes, in particular root

- 16 -

knot nematodes.

The nematocide activity test was performed with larvae of *Meloidogyne incognita* (Kofoid and White) Chitwoody, from soil in the Ferrara area.

5 The test was performed in jars with capacity of 100 g of soil to which second age larvae had been added in order to assess the toxicity of the volatile compounds released by the flour following hydration.

10 The doses equivalent to $20 \cdot 10^{-2}$, $40 \cdot 10^{-2}$ and $120 \cdot 10^{-2}$ Kg/m² of seed flour of *Brassica carinata*, *Barbarea verna* and *Eruca sativa* of example 2 were tested separately. The control, included for assessment of the nematocide activity, consisted of non-treated soil and a fertiliser with two different organic
15 compounds: fowl dung and torrefied leather (applied at the maximum test dose, i.e. $120 \cdot 10^{-2}$ Kg/m²).

20 The test was based on assessment of the mortality of the larvae of *Meloidogyne incognita* in the presence of the flour and the non-biocide flour, with respect to the mortality in untreated soil.

 The table shows the results of the test. The nematocide activity of the flour was expressed as the number of larvae remaining in the soil 10 days after the treatment.

25 The test was performed with four repetitions per treatment, therefore the data shown in table III are

the means of four values.

Table III

Treatment	Dose Kg/m ²	N° larvae 100 g of soil
Blank	-	14
Fowl dung	40.10 ⁻²	8
Leather	40.10 ⁻²	5
<i>B. Carinata</i>	20.10 ⁻²	1
<i>B. Carinata</i>	40.10 ⁻²	0
<i>Eruca sativa</i>	20.10 ⁻²	0
<i>Eruca sativa</i>	40.10 ⁻²	0
<i>B. Verna</i>	20.10 ⁻²	2
<i>B. Verna</i>	40.10 ⁻²	2

The above data indicate that the products released by
 5 the seed flour of *Brassica carinata* A. B., *Eruca sativa*
M. and *B. Verna* following hydration, which activates
 the GLs-MYR system present in it, inhibit to a large
 extent the presence of *Meloidogyne* larvae in the soil.

10 Example 5

This example describes a method for assessing
 insecticide activity, in particular with regard to
 wireworms.

The insecticide activity test was performed with
 15 larvae of *Agriotes sordidus* Illiger, *Agriotes ustulatus*

- 18 -

Schäller, *Agriotes litigiosus* Rossi and *Agriotes brevis* Candeze, produced on breeding farms in Veneto.

The test was performed in jars with capacity of 1.4 l of soil to which larvae of wireworms of the above
5 species at the 6th - 8th stage of development had been added in order to assess the toxicity of the volatile compounds released by the flour following hydration.

The doses equivalent to 20 and 40 quintals ha⁻¹ of seed flour of *Brassica carinata* and *Eruca sativa* of
10 example 2 were tested separately. The control, included for assessment of the insecticide activity, consisted of untreated soil.

The test was based on assessment of the mortality of the larvae of *Agriotes* in the presence of the flour
15 which had undergone hydration, with respect to the mortality in untreated soil. Assessment of mortality was performed one hour after addition of the flour to the soil.

Table II shows the results of the test. The
20 insecticide activity of the flour was expressed as inhibition effectiveness (EI) as a percentage with respect to the control, using the formula: $EI (\%) = [(Mortality\ of\ Control - Mortality\ of\ Treated\ Sample) / Mortality\ of\ Control] \times 100$.

25 The test was performed with four repetitions per treatment, therefore the data shown in table IV are the

means of four values.

Table IV

Treatment with de-oiled flour	EI %
<i>Eruca sativa</i>	92
<i>Brassica carinata</i>	100

The above data indicate that the products released by the seed flour of *Brassica carinata* A. B. and *Eruca sativa* M. following hydration, which activates the GLs-MIR system present in it, inhibit to a large extent the presence of *Agriotes* larvae in the soil.

In particular, the following were observed:

- a high insecticide action of the above-mentioned seed flour in the space of only a few minutes;
- a rapid disappearance of the active compounds (no insecticide activity in the space of only a few days);
- consequent low environmental impact.

No negative interactions were observed.

- 20 -

CLAIMS

1.- Use of a seed flour containing at least one glucosinolate and at least one enzyme chosen from the group consisting of:

- 5 - glucosidasic enzymes, and
 - thioglucosidasic enzymes;
 as soil improver.

2.- Use of a seed flour containing at least one glucosinolate and at least one enzyme chosen from the
10 group consisting of:

- glucosidasic enzymes, and
 - thioglucosidasic enzymes;

 as cytotoxic agent for soil pathogens and soil parasites.

15 3.- Use of a seed flour containing at least one glucosinolate and at least one enzyme chosen from the group consisting of:

- glucosidasic enzymes, and
 - thioglucosidasic enzymes;

20 as fungitoxic agent for soil fungal pathogens.

4.- Use according to claim 2, in which said soil parasites are chosen from the group consisting of:

- nematodes; and
 - wireworms.

25

5.- Use according to claim 2, in which said soil

- 21 -

parasites are chosen from the group consisting of:

- weeds; and
- insects.

6.- Use according to claim 3, in which said flour
5 performs a fungitoxic action towards the fungi of the
genus *Fusarium* and other fungi.

7.- Use according to one of the preceding claims,
in which the seeds are chosen from the group of seeds
consisting of:

- 10 - seeds of *Brassicaceae*;
- seeds of *Capparaceae*;
- seeds of *Koeberliniaceae*;
- seeds of *Moringaceae*;
- seeds of *Resedaceae*; and
- 15 - seeds of *Toviaraceae*.

8.- Use according to one of the preceding claims,
in which the seeds are seeds of *Brassicaceae*.

9.- Use according to one of the claims from 1 to 7,
in which the seeds are seeds of *Capparaceae*.

20 10.- Use according to one of the claims from 1 to
7, in which the seeds have been chosen from the group
consisting of:

- seeds of *Brassica carinata* A. B.;
- seeds of *Eruca sativa* M.; and
- 25 - seeds of *Barbarea verna*.

11.- Use according to one of the preceding claims,

- 22 -

in which said seed flour has been de-oiled.

12.- Use according to claim 11, in which the de-oiling has been performed at a temperature below 75°C.

13.- Use according to one of the previous claims,
5 in which the seeds are of at least two different vegetable varieties.

14.- Use according to one of the preceding claims, in which said enzyme is the enzyme microsinase.

15.- Use according to any one of the preceding
10 claims and comprising an identification phase to identify the seed flour containing at least one glucosinolate and at least one enzyme chosen from the group consisting of:

- glucosidasic enzymes, and
- 15 - thioglucosidasic enzymes.

16.- Use according to claim 2, in which said soil parasites are chosen from the group consisting of:

- *Fusarium culmorum*;
- *Meloidogyne incognita* (Kofoid and White)
- 20 *Chitwoody*;
- *Agriotes sordidus*;
- *Agriotes utulatus*; and
- *Agriotes brevis*.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IT 03/00514

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N65/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, PAJ, EPO-Internal, WPI Data, CAB Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1996 ELBERSON LESLIE R ET AL: "Toxicity of rapeseed meal-amended soil to wireworms, Limonius californicus (Coleoptera: Elateridae)" Database accession no. PREV199799288779 XP002265105 abstract & JOURNAL OF AGRICULTURAL ENTOMOLOGY, vol. 13, no. 4, 1996, pages 323-330, ISSN: 0735-939X</p> <p style="text-align: center;">--- -/--</p>	<p>1,2,4,5, 7,8,14, 15</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

16 December 2003

Date of mailing of the international search report

13/01/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lamers, W

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/IT 03/00514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; June 2001 (2001-06) HAFEZ S L ET AL: "Efficacy of oil seed meals for management of Heterodera schachtii and Meloidogyne chitwoodi under green house conditions" Database accession no. PREV200100448302 XP002265106 abstract & PHYTOPATHOLOGY, vol. 91, no. 6 Supplement, June 2001 (2001-06), page S135 Joint Meeting of the American Phytopathological Society, the Mycological Society of America, and the; Salt Lake City, Utah, USA; August 25-29, 2001 ISSN: 0031-949X</p> <p>---</p>	1,2,4,7, 8,15,16
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1997 BOREK VLADIMIR ET AL: "Toxicity of rapeseed meal and methyl isothiocyanate to larvae of the black vine weevil (Coleoptera: Curculionidae)" Database accession no. PREV199799422007 XP002265107 abstract & JOURNAL OF ECONOMIC ENTOMOLOGY, vol. 90, no. 1, 1997, pages 109-112, ISSN: 0022-0493</p> <p>---</p>	1,2,5,7, 8,14,15
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; December 1997 (1997-12) SIDDIQUI M A ET AL: "Integrated control of plant parasitic nematodes with organic soil amendments /nematicides and ploughing on okra" Database accession no. PREV199900282951 XP002265108 abstract & PROCEEDINGS OF THE INDIAN NATIONAL SCIENCE ACADEMY PART B BIOLOGICAL, vol. 63, no. 6, December 1997 (1997-12), pages 545-550, Dec., 1997 ISSN: 0073-6600</p> <p>---</p> <p style="text-align: center;">-/--</p>	1,2,4,7, 8,10-12, 14,15

INTERNATIONAL SEARCH REPORT

Internz Application No
PCT/IT 03/00514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB; S.ANVER ET AL.: "Organic management of concomitant Meloidogyne incognita and Rotylenchus reniformis on chickpea" retrieved from STN-INTERNATIONAL Database accession no. 2000:46428 CABA XP002265109 abstract & ALLELOPATHY JOURNAL, vol. 7, no. 1, 2000, pages 79-84.</p>	1,2,4,7, 8,10-16
X	<p>DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB; S.ANVER ET AL.: "Biological control of soil nematodes associated with linseed" retrieved from STN-INTERNATIONAL Database accession no. 2002:79103 CABA XP002265110 abstract & ARCHIVES OF PHYTOPATHOLOGY AND PLANT PROTECTION, vol. 34, no. 2, 2001, pages 101-109,</p>	1,2,4,7, 8,10-15
X	<p>DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB; S.A.TIYAGI ET AL.: "Biodegradable effects of oilseed cakes on plant parasitic nematodes and soil-inhabiting fungi infesting Trigonella foenum-graecum and Phaseolus aureus" retrieved from STN-INTERNATIONAL Database accession no. 2002:186203 CABA XP002265111 abstract & INDIAN JOURNAL OF NEMATOLOGY, vol. 32, no. 1, 2002, pages 47-57,</p>	1-4,6-8, 10-16
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1995 TIYAGI SARTAJ A ET AL: "Efficacy of oil-seed cakes against plant-parasitic nematodes and soil-inhabiting fungi on mungbean and chickpea" Database accession no. PREV199598265524 XP002265112 abstract & BIORESOURCE TECHNOLOGY, vol. 51, no. 2-3, 1995, pages 233-239, ISSN: 0960-8524</p>	1-8, 10-16

-/--

INTERNATIONAL SEARCH REPORT

Intern~~e~~ Application No
PCT/IT 03/00514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; April 1998 (1998-04) SARWAR M ET AL: "Biofumigation potential of brassicas. III. In vitro toxicity of isothiocyanates to soil-borne fungal pathogens" Database accession no. PREV199800486065 XP002265113 abstract & PLANT AND SOIL, vol. 201, no. 1, April 1998 (1998-04), pages 103-112, ISSN: 0032-079X</p>	1-3,6-8, 14,15
X	<p>--- DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; May 2000 (2000-05) TSAO RONG ET AL: "Factors affecting the dissolution and degradation of oriental mustard-derived sinigrin and allyl isothiocyanate in aqueous media" Database accession no. PREV200000301573 XP002265114 abstract & JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY, vol. 48, no. 5, May 2000 (2000-05), pages 1898-1902, ISSN: 0021-8561</p>	1,2,4,7, 8,14,15
X	<p>--- DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1993 KUMAR KRISHAN ET AL: "Efficacy of organic-amendments on the yield and control of Aphelenchoides composticola infesting mushroom, Agaricus bisporus (Lange) Sing" Database accession no. PREV199497502483 XP002265115 abstract & INDIAN JOURNAL OF NEMATOLOGY, vol. 23, no. 1, 1993, pages 82-86, ISSN: 0303-6960</p> <p>--- -/--</p>	1,2,4,7, 8,11-15

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IT 03/00514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB; S.S.JAKHAR ET L.: "Effect of soil amendment with some oil cakes on root rot of cotton caused by Rhizoctania sp." retrieved from STN-INTERNATIONAL Database accession no. 2003:7637 CABA XP002265116 abstract & PLANT DISEASE RESEARCH, vol. 17, no. 1, 2002, pages 16-20,</p>	1-3,6-8, 10-15
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; July 2001 (2001-07) MAZZOLA MARK ET AL: "Suppression of specific apple root pathogens by Brassica napus seed meal amendment regardless of glucosinolate content" Database accession no. PREV200100391628 XP002265117 abstract & PHYTOPATHOLOGY, vol. 91, no. 7, July 2001 (2001-07), pages 673-679, ISSN: 0031-949X</p>	1,2,4,7, 8,14,15
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1995 BROWN PAUL D ET AL: "Glucosinolate-containing plant tissues as bioherbicides" Database accession no. PREV199698721413 XP002265118 abstract & JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY, vol. 43, no. 12, 1995, pages 3070-3074, ISSN: 0021-8561</p>	1,2,5,7, 8,11-15
P,X	<p>US 6 596 323 B1 (DEUEL WES A) 22 July 2003 (2003-07-22) the whole document</p>	1


-/--

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/IT 03/00514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1997 MANICI LUISA M ET AL: "In vitro fungitoxic activity of some glucosinolates and their enzyme-derived products toward plant pathogenic fungi" Database accession no. PREV199799697211 XP002265119 abstract & JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY, vol. 45, no. 7, 1997, pages 2768-2773, ISSN: 0021-8561</p>	1-16
A	<p>--- S.PALMIERI ET AL.: "The Glucosinolate Myronase System - A Natural & Practical Tool for Biofumigation" HORTICULTURE BIOFUMIGATION UPDATE, no. 12, November 2000 (2000-11), pages 1-2, XP002265103 the whole document</p>	1-16
A	<p>--- DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1998 ANGELINI L ET AL: "Antigerminative activity of three glucosinolate-derived products generated by myrosinase hydrolysis" Database accession no. PREV199900264863 XP002265120 abstract & SEED SCIENCE AND TECHNOLOGY, vol. 26, no. 3, 1998, pages 771-780, ISSN: 0251-0952</p>	1-16
A	<p>--- DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1991 RODMAN J E: "A TAXONOMIC ANALYSIS OF GLUCOSINOLATE-PRODUCING PLANTS PART 2 CLADISTICS" Database accession no. PREV199293025043 XP002265121 abstract & SYSTEMATIC BOTANY, vol. 16, no. 4, 1991, pages 619-629, ISSN: 0363-6445</p> <p>--- -/--</p>	1-16

INTERNATIONAL SEARCH REPORT

Inter  Application No
PCT/IT 03/00514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	D.FROHNE ET AL.: "Systematik des Pflanzenreiches" 1998 , WISSENSCHAFTLICHE VERLAGSGESELLSCHAFT MBH , STUTTGART, DE XP002265104 Ordnungsgruppe Senfartige page 204 -page 211 ---	1-16
A	WO 97 14309 A (CANADA MAJESTY IN RIGHT OF) 24 April 1997 (1997-04-24) examples 1,2 -----	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IT 03/00514

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6596323	B1	22-07-2003	NONE
WO 9714309	A	24-04-1997	US 5747056 A 05-05-1998
		AT 200181 T 15-04-2001	
		AU 7081696 A 07-05-1997	
		CA 2233602 A1 24-04-1997	
		WO 9714309 A1 24-04-1997	
		DE 69612391 D1 10-05-2001	
		DE 69612391 T2 31-10-2001	
		EP 0855860 A1 05-08-1998	
		ZA 9608648 A 11-07-1997	